

Our examination of a binary star mass distribution [1, 2] shows that cores of these stars in their stable state should be composed from plasma consisting of electrons and neutron-excess nuclei (the same as 3H , 4H , 5H and 6He , 8He , ^{10}He). These nuclei are β -active under "normal" terrestrial conditions. They get stable in central regions of stars under pressure of plasma electron gas. This pressure averaged over a plasma cell at plasma density about 10^{25} particule/ cm^3 and temperature about 10^7K [1, 2] amounts to 10^6GPa . It can be considered as the lowest bond of the electron pressure acting on nuclei. As the density of electron gas in the nearest vicinity of a nucleus is much more than the density averaged over a cell volume, its pressure on nucleus is substantially higher than the mentioned value. It is impossible to achieve Such a pressure at laboratory conditions, as well as to influence the rate of β -decay. Let us estimate how much can the electron gas slow down the decay rate at laboratory conditions. The most suitable object for this investigation is tritium. It is subjected to β -decay with a highest possible energy of an outgoing electron equal to $E_\beta=18.6$ kev. The ionization energy of hydrogen is $E_i=13.6$ ev. The energy of this level can be held by electrons in plasma, which is created by a gas discharge in tritium gas. If nuclei experiencing β -decay compose atomic substance, they emit a full spectrum of electron energy from 0 to E_β . If these nuclei compose an electron-nuclear plasma, the outgoing of decay electrons with energies less than E_i must be limited by the Pauli principle. The part of a phase volume of this state is

$$\delta = \left(\frac{E_i}{E_\beta} \right)^{3/2} \approx 10^{-4.5} \quad (1)$$

The probability of a electron outgoing with energy close to the threshold E_β is small in comparison to the probability of a electron outgoing with a small energy $E \ll E_\beta$. It must several (approximately 3) times increase the effect of plasma influence and we can expect that the influence of plasma on β -decay reaches a value approximately equal to 10^{-4} . To have the statistically valid measured result, the total amount N of decays during the time of measurement must be at the level

$$N > \left(\frac{3}{\delta} \right)^2 \approx 10^8, \quad (2)$$

It seems quite realistic to achieve this effect with enough tritium in the experimental device and a reasonable rate and time of the measurement.

Список литературы

- [1] Vasiliev B.V. - Nuovo Cimento B, 2001, v.116, pp.617-634.
- [2] Vasiliev B.V. - astro-ph/0512468